


# **Safely Reducing Primary Cesarean Section Utilizing the PROVIDE Toolbox in a Faith-Based Hospital**

**Minyon Outlaw, DNP, CNM, WHNP-BC**




# Specific Aim


- Safely reduce primary cesareans
- Population: nulliparous, term, singleton, vertex (NTSV)
- Decrease rate from 29.1% to 24% in a 90-day period




## NTSV C/S Rate: Birth Log and EHR Comparison



## Labor Dystocia Checklist Score



## Team Perceptions & Confidence Survey



# Discussion


---

## Strengths of project

- Team Spirit
- Cost effective
- Leadership Support
- “Doula the Nurse”

## Summary of key findings


- Reduction of NTSV C/S from 29.1% to **20.2%**
- Labor dystocia score improved from 26% to **56.4%**
- Team engagement mean score increased from 3.2 to **4.2**



(AHRQ, 2018; Wise and Jolles, 2019)



# Conclusion





# References

Agency for Healthcare Research and Quality. (2018). *TeamSTEPPS pocket guide*. Retrieved from <https://www.ahrq.gov/teamstepps/instructor/essentials/pocketguideapp.html>

Caughey, A. B., Cahill, A. G., Guise, J., & Rouse, D. J. (2014). Safe prevention of the primary cesarean delivery. *American Journal of Obstetrics and Gynecology*, 210(3), 179-193. doi:10.1016/j.ajog.2014.01.026

Florida Perinatal Quality Collaborative, [FPQC]. *Promoting primary vaginal deliveries (PROVIDE) tool box for hospital implementation*. Retrieved from <https://health.usf.edu/publichealth/chiles/fpqc/provide/toolbox>

HealthyPeople2020. (2019). *Reduce cesarean births among low-risk women with no prior cesarean births*. Retrieved from [https://www.healthypeople.gov/node/4900/data\\_details](https://www.healthypeople.gov/node/4900/data_details)

Ogrinc, G. S., Headrick, L. A., Barton, A. J., Dolansky, M. A., Madigosky, W. S., and Miltner, R. S. (3<sup>rd</sup> ed.). (2018). *Fundamentals of health care improvement: A guide to improving your patients' care*. Boston, MA. The Joint Commission Resources

Sappenfield, W. M. (2019). *Background and significance of cesareans*. Retrieved from <https://health.usf.edu/-/media/Files/Public%20Health/Chiles%20Center/FPQC/PROVIDEBackground%20Slides%20-%Sappenfield.ashx?1a=en&hash=AE3A0390F824319B33E7EE2FBB8E01348BC68AAA>

Smith, H., Peterson, N., Lagrew, D., and Main, E. (2017). *Toolkit to support vaginal birth and reduce primary cesarean sections*. Retrieved from file:///C:/Users/minyo/Downloads/VBirth%20Toolkit%20Final\_V09.20.17.

The Joint Commission. (2019). *Leading hospital improvement*. Retrieved from [https://www.jointcommission.org/the\\_view\\_from\\_the\\_joint\\_commission/public\\_reporting\\_of\\_high\\_cesarean\\_rates\\_to\\_begin\\_in\\_july\\_2020/](https://www.jointcommission.org/the_view_from_the_joint_commission/public_reporting_of_high_cesarean_rates_to_begin_in_july_2020/)

Wise, G. and Jolles, D. (2019). Promoting effective care: Reducing primary cesarean births through team engagement and standardization of care at a community hospital. *Nurs Forum*. DOI: <https://doi.org/10.1111/nuf.1238>